

UHF flows and the flip automorphism

A. Kishimoto

Department of Mathematics, Hokkaido University, Sapporo 060, Japan

November 2000

1 Introduction

By a *flow* on a unital C^* -algebra A we mean a strongly continuous one-parameter automorphism group. The infinitesimal generator δ_α of a flow α is a closed derivation in A , by which we mean that δ_α is a closed linear operator which is defined on a dense $*$ -algebra $D(\delta_\alpha)$ and satisfies that $\delta_\alpha(x)^* = \delta_\alpha(x^*)$ and $\delta_\alpha(xy) = \delta_\alpha(x)y + x\delta_\alpha(y)$. See [1, 4, 17] for the general theory of derivations.

If α is a flow and $u = (u_t)$ is a one-parameter family of unitaries of A such that $t \mapsto u_t$ is continuous and $u_s\alpha_s(u_t) = u_{s+t}$ for all $s, t \in \mathbf{R}$, we say that u is an α -cocycle. Then $\text{Ad } u\alpha : t \mapsto \text{Ad } u\alpha_t$ is a flow and is called a cocycle perturbation of α . If u is differentiable with

$$h = -i \frac{du_t}{dt} \Big|_{t=0},$$

then $\text{Ad } u\alpha$ is an inner perturbation of α in the sense that $\delta_{\text{Ad } u\alpha} = \delta_\alpha + \text{ad } ih$, where $\text{ad } ih$ is the inner derivation defined by $\text{ad } ih(x) = i(hx - xh)$ for $x \in A$. We write $\text{Ad } u\alpha$ also as $\alpha^{(h)}$ in this case.

If v is a unitary which is not in the domain $D(\delta_\alpha)$, then the flow

$$\text{Ad } v\alpha \text{ Ad } v^* : t \mapsto \text{Ad } v\alpha_t(v^*)\alpha_t,$$

which is conjugate to α , is a cocycle perturbation of α but not an inner perturbation. In general a cocycle perturbation is conjugate to an inner perturbation since any α -cocycle u is cohomologous to a differentiable one w , i.e., $u_t = vw_t\alpha_t(v^*)$ for some unitary v (see [12]).

By an inner flow we mean a flow given by $t \mapsto \text{Ad } e^{iht} = e^{t \text{ad } ih}$ for some self-adjoint element h in A . We say that α is approximately inner if there is a sequence (h_n) in A_{sa} such that $\alpha_t = \lim \text{Ad } e^{ihn_t}$, i.e., $\alpha_t(x) = \lim \text{Ad } e^{ihn_t}(x)$ for every $t \in \mathbf{R}$ and $x \in A$, or equivalently, uniformly continuous in t on every compact subset of $\mathbf{R}$ and every $x \in A$.

When A is an AF C^* -algebra, we call α an *AF flow* if there is an increasing sequence (A_n) of finite-dimensional C^* -subalgebras of A such that $1_A \in A_1$, $A = \overline{\bigcup_n A_n}$ and $\alpha_t(A_n) = A_n$ for all n and t . In this case there is an h_n in $(A_n)_{sa}$ such that

$\alpha_t|_{A_n} = \text{Ad } e^{ih_n t}|_{A_n}$ and hence α is approximately inner. (The term AF flow is coined in [2] while the adjective *locally representable* is used in [12] and elsewhere to refer to the same object.)

When A is a UHF C^* -algebra, we call α a *UHF flow* if there is an increasing sequence (A_n) of full matrix C^* -subalgebras of A such that $1_A \in A_1$, $A = \overline{\cup_n A_n}$, and $\alpha_t(A_n) = A_n$ for all n and t . If we set $B_n = A_n \cap A'_{n-1}$ with $A_0 = 0$, then $A \cong \bigotimes_1^\infty B_n$ and $\alpha_t(B_n) = B_n$ for all n . Namely α is of infinite tensor product type. Note that UHF flows are AF flows and there are AF flows on UHF C^* -algebras which are not UHF flows. This follows because there are AF flows which have more than one KMS states for some temperature while UHF flows always have a unique KMS state for any temperature (see e.g., [12]).

We call α a *compact flow* if the closure of $\alpha_{\mathbf{R}}$ in $\text{Aut}(A)$, the automorphism group of A , is compact, or equivalently, if there is an increasing sequence (V_n) of finite-dimensional subspaces of A such that $A = \overline{\cup_n V_n}$ and $\alpha_t(V_n) = V_n$ for all n and t . A periodic flow is compact but there are more. If α is an AF flow, then it is compact.

In [13] we refer to locally inner (or locally representable) flows as a generalization of AF flows. That is, a flow α is *locally inner* if there is an increasing sequence (A_n) of (arbitrary) C^* -subalgebras of A such that $A = \overline{\cup_n A_n}$ and α leaves A_n invariant and restricts to an inner flow on A_n . Apparently AF flows are locally inner and locally inner flows are approximately inner. But there seems to be no obvious relation between compact flows and locally inner flows.

When A is a UHF C^* -algebra, we have the following implications for flows:

$$\text{UHF flows} \Rightarrow \text{AF flows} \begin{cases} \Rightarrow & \text{Locally inner flows} \\ \Rightarrow & \text{Compact flows} \end{cases}$$

where the reverse implications are false (for the latter note that the examples of non-AF flows constructed in [13] are locally inner and compact).

There is certainly a non-compact flow (e.g., a flow with some asymptotic abelianess), but we do not seem to know if there is a flow which is not even a cocycle perturbation of a compact flow. If there is a flow which is not approximately inner (against the Powers-Sakai Conjecture [17]), it is likely that we have such a one among the compact (or even periodic) flows since there is such a one for some simple AF C^* -algebra, which is obtained from [14] and Lin's classification theorem for simple C^* -algebras of tracial topological rank zero. Hence there may be no inclusion relation between the compact flows and the approximately inner flows.

In this note we will briefly discuss compact flows and then look into UHF flows, a simplest kind of flows!

We will show in 2.2 that if α is an approximately inner compact flow on A , then the domain $D(\delta_\alpha)$ of the generator δ_α contains a maximal abelian C^* -subalgebra (masa for short) of A , a property which obviously holds for locally inner flows. We here refer to a result that the AF flows are characterized by the property that the domain contains a canonical AF masa (see [13] for details).

When A is a UHF C^* -algebra such that $A \otimes A \cong A$, we call a UHF flow γ on A *universal* if $\gamma \otimes \alpha$ is cocycle conjugate to γ for any UHF flow α on A (i.e., there is an isomorphism φ of $A \otimes A$ onto A such that $\gamma \otimes \alpha$ is a cocycle perturbation of $\varphi^{-1}\gamma\varphi$). There exist universal UHF flows on A (as shown in 3.1) and they are mutually cocycle conjugate, or even *almost conjugate* (see 4.5). In the case of the CAR algebras we will construct a universal UHF flow in a simple way extending a result given in [12]; see 5.1.

When α is a flow on A and σ is the flip automorphism of $A \otimes A$, i.e., $\sigma(x \otimes y) = y \otimes x$, $x, y \in A$, we say that σ is α -*invariantly approximately inner* if there is a sequence (u_n) of unitaries in $A \otimes A$ such that $\sigma = \lim \text{Ad } u_n$ and $\|(\alpha_t \otimes \alpha_t)(u_n) - u_n\| \rightarrow 0$ for each $t \in \mathbf{R}$, or equivalently uniformly in t on every compact subset of $\mathbf{R}$. We shall prove for an approximately inner flow α that $\alpha \otimes \gamma$ is a cocycle perturbation of a (universal) UHF flow if and only if σ is α -invariantly approximately inner, where γ is a universal UHF flow; see 4.6. We do not really have any application of this result (except for a result on quasi-UHF flows; see 4.7) but this is a first attempt to characterize UHF flows; see [13, 2] for some results on AF flows. We shall also note that if the flip is α -smoothly approximately inner, i.e., if we replace the condition $\|(\alpha_t \otimes \alpha_t)(u_n) - u_n\| \rightarrow 0$ by that $(\alpha_t \otimes \alpha_t)(u_n)$ is equi-continuous in the above definition, then α has a unique KMS state for each inverse temperature, see 4.3. This follows from [8] and applies to the one-dimensional quantum lattice systems, thus providing another proof of the well-known uniqueness result.

2 Compact flows

Proposition 2.1 *Let A be a separable C^* -algebra and α a flow on A . Then the following conditions are equivalent:*

1. *The closure of $\{\alpha_t \mid t \in \mathbf{R}\}$ in $\text{Aut}(A)$ is compact.*
2. *For each $x \in A$ the closure of $\{\alpha_t(x) \mid t \in \mathbf{R}\}$ is compact.*
3. *For each x of a dense subset of A the closure of $\{\alpha_t(x) \mid t \in \mathbf{R}\}$ is compact.*
4. *The linear span of $A_p = \{x \in A \mid \alpha_t(x) = e^{ipt}x\}$ for all $p \in \mathbf{R}$ is dense in A .*
5. *There is an increasing sequence (V_n) of finite-dimensional subspaces of A such that $A = \overline{\bigcup_n V_n}$ and $\alpha_t(V_n) = V_n$ for all n and t .*

Proof. (1) $\Rightarrow$ (2) $\Rightarrow$ (3) is obvious. To show that (3) $\Rightarrow$ (1) let (x_n) be a dense sequence (x_n) in the dense subset of A given in (3) and let (t_n) be a sequence in $\mathbf{R}$; then there is a subsequence $(t_{n(k)})$ such that both $(\alpha_{t_{n(k)}}(x_m))$ and $(\alpha_{-t_{n(k)}}(x_m))$ converge for any m . Then the (strong) limit of $(\alpha_{t_{n(k)}})$ exists as an automorphism. This implies (1).

If (1) holds, then the closure of $\alpha_{\mathbf{R}}$ is a compact abelian group and hence (4) follows. It is immediate that (4) $\Rightarrow$ (5) $\Rightarrow$ (3).

We recall that α is said to be a compact flow if the conditions in 2.1 are satisfied.

Proposition 2.2 *Let A be a unital separable simple C^* -algebra and α a flow on A . If α is an approximately inner compact flow, then $D(\delta_\alpha)$ contains a maximal abelian C^* -subalgebra of A .*

Proof. Since α is approximately inner, α has a pure ground state ω . If G denotes the closure of $\alpha_{\mathbf{R}}$, then ω is left invariant under G . Thus the GNS representation associated with ω is a G -covariant irreducible representation π of A . Since A is simple, π is faithful.

Let U be a continuous representation of G on $\mathcal{H}_\pi$ such that $U_g \pi(x) U_g^* = \pi g(x)$ for $x \in A$. There is an orthogonal family $(E_p)_{p \in \hat{G}}$ of projections such that $U_g = \sum_p \langle g, p \rangle E_p$, where $\hat{G}$ is the character group of G ; $\hat{G}$ is a countable discrete abelian group. Let $A^G = \{x \in A \mid \forall g \in G, g(x) = x\}$ and denote by π_p the representation of A^G on $E_p \mathcal{H}_\pi$, i.e., $\pi_p(a) = \pi(a)|_{E_p \mathcal{H}_\pi}$, $a \in A^G$. It follows that $\{\pi_p \mid p \in \hat{G}\}$ is a disjoint family of irreducible representations of A^G since $\pi(A^G)' = U_G''$. Under this circumstance we will show that there is a maximal abelian C^* -subalgebra (masa) of A^G which is also a masa in A .

By the following lemma there is an $h \in (A^G)_{sa}$ such that $\pi(h) \cong \oplus_{p \in \hat{G}} \pi_p(h)$ is diagonal and all the eigenvalues of $\pi(h)$ have multiplicity one. Let C be a masa of A^G such that $C \ni h$. Since $A \cap C'$ is G -invariant, $A \cap C'$ is the closed linear span of the eigenspaces of $G|_{A \cap C'}$. Let $x \in A \cap C'$ be a nonzero element such that $g(x) = \langle g, q \rangle x$, $g \in G$ for some $q \in \hat{G}$. Then there must be a $p \in \hat{G}$ and a unit vector ξ in $E_p \mathcal{H}$ such that $\pi(x)\xi \neq 0$ and $\pi(h)\xi = \lambda\xi$ for some $\lambda \in \mathbf{R}$. Since $\pi(x)\xi \in E_{p+q} \mathcal{H}$ and $\pi(h)\pi(x)\xi = \pi(x)\pi(h)\xi = \lambda\pi(x)\xi$, $\pi(x)\xi$ is a constant multiple of ξ . This implies that $q = 0$ and then $x \in C$. Thus we can conclude that C is a masa in A .

Lemma 2.3 *Let A be a separable C^* -algebra and let (π_n) be a sequence of irreducible representations of A such that if $m \neq n$ then π_m is disjoint from π_n . If (S_n) is a sequence of dense subsets of $\mathbf{R}$ then there is an $h \in A_{sa}$ such that $\pi_n(h)$ is diagonal and $\text{PSp}(\pi_n(h)) \subset S_n$ for all n . Furthermore h can be chosen so that all the eigenvalues of $\pi_n(h)$ have multiplicity one for every n .*

Proof. We will construct a sequence (h_n) in A_{sa} and an increasing sequence $(e_{kn})_{n \geq k}$ of finite-dimensional projections on $\mathcal{H}_k = \mathcal{H}_{\pi_k}$ for each k such that $\|h_n\| < 2^{-n}$, $\lim_n e_{kn} = 1$, $\pi_k(h_n)e_{k,n-1} = 0$ for $k = 1, 2, \dots, n-1$, and for $H_n = h_1 + h_2 + \dots + h_n$,

$$\begin{aligned} [\pi_k(H_n), e_{kn}] &= 0, \quad k = 1, 2, \dots, n \\ \text{PSp}(\pi_k(H_n)|_{e_{kn} \mathcal{H}_k}) &\subset S_k, \quad k = 1, 2, \dots, n, \end{aligned}$$

where PSp denotes the set of eigenvalues. Then we will let $h = \lim_n H_n$ and then since $\pi_k(h)e_{k,n} = \pi_k(H_n)e_{k,n}$, we will have that $\text{PSp}(\pi_k(h)) \subset S_k$ for all k .

To construct such sequences as above, we will argue inductively by using Kadison's transitivity theorem (see, e.g., 1.21.16 of [18]). Suppose that we have constructed $h_1, h_2, \dots, h_n$ and e_{km} with $n \geq m \geq k$ satisfying the above conditions. With $H_n =$

$h_1 + \cdots + h_n$ let E_k be the spectral measure of $\pi_k(H_n)$ for $k = 1, 2, \dots, n+1$. For each k we find a finite family F_k of disjoint translates of $[0, 2^{-n-1})$ and a family $\{\xi(I)\}_{I \in F_k}$ of unit vectors in $(1 - e_{kn})\mathcal{H}_k$ such that $E_k(I)\xi(I) = \xi(I)$ and the linear span of $\xi(I)$, $I \in F_k$ is so large that it almost contains any prescribed vector from $(1 - e_{kn})\mathcal{H}_k$. Choose $\lambda_I \in S_k \cap I$ and let P_I denote the projection onto the linear space spanned by $\xi(I)$ and $(\pi_k(H_n) - \lambda_I)\xi(I)$. Then the family $(P_I)_{I \in F_k}$ is mutually orthogonal and the sum $P_k = \sum_I P_I$ is orthogonal to e_{kn} . It follows that $\|P_k(\pi_k(H_n) - \sum_I \lambda_I P_I)P_k\| = \|\sum_I P_I(\pi_k(H_n) - \lambda_I P_I)P_I\| < 2^{-n-1}$. We will then choose an $h_{n+1} \in A_{sa}$ such that $\|h_{n+1}\| \leq 2^{-n-1}$, $[\pi_k(h_{n+1}), P_k] = 0$ for $k \leq n+1$, $\pi_k(h_{n+1})e_{kn} = 0$ for $k \leq n$, and $\pi_k(h_{n+1})P_k = -(\pi_k(H_n) - \sum_I \lambda_I P_I)P_k$ for $k = 1, 2, \dots, n+1$. Then we obtain that $\pi_k(H_n + h_{n+1})\xi(I) = \lambda(I)\xi(I)$. We let $e_{k,n+1}$ be the sum of e_{kn} (0 if $k = n+1$) and the projection onto the linear span of $\xi(I)$, $I \in F_k$. Thus we have constructed h_{n+1} , $e_{k,n+1}$ as required. (A similar argument is used in [9].)

Remark 2.4 In 2.2 the assumption that α is approximately inner is made to ensure that there is a G -covariant irreducible representation for $G = \overline{\alpha_{\mathbf{R}}}$. This assumption certainly is not necessary as we can see from the following example. If α is a flow on the Cuntz algebra $\mathcal{O}_2 = C^*(s_1, s_2)$ such that $\alpha_t(s_j) = e^{i\mu_j t} s_j$, where μ_1, μ_2 are rationally independent, then α is compact and G is the gauge action γ of $\mathbf{T}^2$ given by $\gamma_{z_1, z_2}(s_j) = z_j s_j$. In this case α is not approximately inner but γ has a covariant irreducible representation. Hence the conclusion of 2.2 follows for this α (though it is easy to show that directly). On the other hand there is an example of a compact flow where G does not have a covariant irreducible representation. For example if α is a flow on the irrational rotation C^* -algebra $A_\theta = C^*(u_1, u_2)$ with $u_1 u_2 = e^{2\pi i \theta} u_2 u_1$ and θ irrational such that $\alpha_t(u_j) = e^{i\mu_j t} u_j$, where μ_1, μ_2 are rationally independent, then G is the gauge action γ of $\mathbf{T}^2$ given by $\gamma_{z_1, z_2}(u_j) = z_j u_j$. Since γ is ergodic, only the tracial representation is covariant though there is an α -covariant irreducible representation [9]. In this case we do not know whether the domain $D(\delta_\alpha)$ contains a masa or not.

3 Universal UHF flows

We say that a flow α on A is *almost conjugate* to a flow β on B and denote it by $\alpha \stackrel{ac}{\sim} \beta$ if for any $\epsilon > 0$ there is an isomorphism φ of A onto B such that $\|\alpha_t - \varphi^{-1}\beta_t\varphi\| < \epsilon$ for $t \in [-1, 1]$. If A (or B) is separable and simple and $\alpha \stackrel{ac}{\sim} \beta$, then $\alpha \stackrel{cc}{\sim} \beta$, i.e., α is cocycle conjugate to β (see 1.2 of [12]). To prove this we use the fact that if A is simple and $\|\alpha_t - \varphi^{-1}\beta_t\varphi\| < 2$, then there is a unitary $u_t \in A$ such that $\alpha_t = \text{Ad } u_t \varphi^{-1}\beta_t\varphi$. To get an α -cocycle from the family (u_t) of unitaries so obtained for small $|t|$, we may use 8.1 of [16], which requires separability.

Let A be a UHF C^* -algebra such that $A \otimes A \cong A$. We recall that a flow γ on A is a universal UHF flow if $\gamma \otimes \alpha \stackrel{cc}{\sim} \gamma$ for any UHF flow α on A .

Proposition 3.1 *If A is a UHF C^* -algebra such that $A \otimes A \cong A$, then there is a universal UHF flow γ on A .*

Proof. There exists a (finite or infinite) sequence (p_i) of prime numbers such that $A \cong \otimes_i M_{p_i}^\infty$. Let S be the set of integers which are of the form $p_1^{k_1} p_2^{k_2} \cdots p_n^{k_n}$ with $k_i \geq 0$. Then it follows that $A \cong \otimes_{q \in S} M_{q^\infty}$. For each $q \in S$ let (h_n) be a dense sequence in the self-adjoint diagonal matrices of M_q . We define a flow $\gamma^{(q)}$ on the UHF C^* -algebra M_{q^∞} by

$$\bigotimes_{n=1}^{\infty} \text{Ad } e^{ith_n}$$

and define a flow γ on A by $\otimes_{q \in S} \gamma^{(q)}$.

If α is a UHF flow on A , then there are an infinite sequence (q_n) in S and a sequence (k_n) with $k_n \in M_{q_n}$ being self-adjoint and diagonal such that α is conjugate to

$$\bigotimes_{n=1}^{\infty} \text{Ad } e^{itk_n}.$$

Hence $\gamma \otimes \alpha$ is of the same form as $\gamma = \otimes_{q \in S} \gamma^{(q)}$ (by tensoring $\gamma^{(q)}$ with $\otimes_{n: q_n=q} \text{Ad } e^{itk_n}$) but the sequence (h'_n) defining the flow on M_{q^∞} may be different from the (h_n) defining $\gamma^{(q)}$ for some $q \in S$. But since they are dense in the self-adjoint diagonal matrices of M_{q^∞} for any $q \in S$, we can conclude that $\gamma \otimes \alpha$ is cocycle conjugate to γ , i.e., $\gamma \otimes \alpha \overset{cc}{\sim} \gamma$. (As a matter of fact $\gamma \otimes \alpha \overset{ac}{\sim} \gamma$.)

Let $F(A)$ denote the set of flows on A . We give a topology on $F(A)$ by

$$d(\alpha, \beta) = \sum_{n=1}^{\infty} 2^{-n} \max_{|t| \leq 1} \|\alpha_t(x_n) - \beta_t(x_n)\|,$$

where (x_n) is a dense sequence of the unit ball of A . That is, α_n converges to α if $\alpha_{nt}(x)$ converges to $\alpha_t(x)$ uniformly in t on $[-1, 1]$ for all $x \in A$. Let $AIF(A)$ denote the set of approximately inner flows on A , i.e., the closure of inner flows.

Proposition 3.2 *Let A be a UHF C^* -algebra such that $A \otimes A \cong A$ and γ a universal UHF flow on A . Then*

$$\{\varphi^{-1} \gamma \varphi \mid \varphi \in \text{Aut}(A)\} \equiv C(\gamma)$$

is dense in $AIF(A)$.

Proof. Let $h \in A_{sa}$. It suffices to show that the closure of $C(\gamma)$ contains the inner flow $t \mapsto \text{Ad } e^{ith}$.

Since γ is a universal UHF flow, it follows that $\gamma \otimes (\otimes^\infty \text{Ad } e^{ith}) \overset{cc}{\sim} \gamma$, i.e., there is an isomorphism φ of $A \otimes (\otimes^\infty A) \equiv \otimes^\infty A$ onto A and a $k \in A_{sa}$ such that

$$\gamma_t \otimes \left(\bigotimes_{n=1}^{\infty} \text{Ad } e^{ith} \right) = \varphi^{-1} \gamma_t^{(k)} \varphi,$$

where $\gamma^{(k)}$ is the inner perturbation of γ by $\text{ad } ik$. Let φ_n denote the homomorphism of A into A obtained by restricting φ to the $n+1$ 'st factor of $\otimes^\infty A$. Then we have that

$$\lim_{k \rightarrow \infty} (\varphi_n \text{Ad } e^{ith}(x) - \gamma_t \varphi_n(x)) = 0$$

uniformly in t on $[-1, 1]$ for every $x \in A$. Since there are isomorphisms $\tilde{\varphi}_n$ of A onto A such that $\lim(\varphi_n(x) - \tilde{\varphi}_n(x)) = 0$ for every $x \in A$, we may assume that φ_n 's are all isomorphisms of A onto A and obtain that $\text{Ad } e^{ith}$ is the limit of $\varphi_n^{-1}\gamma\varphi_n \in C(\gamma)$.

Remark 3.3 If a flow α absorbs the universal UHF flow γ , i.e., $\alpha \otimes \gamma \stackrel{ec}{\sim} \alpha$, then the closure of $C(\alpha)$ contains $AIF(A)$. This follows since for $h \in A_{sa}$,

$$\alpha \otimes \left(\bigotimes_{n=1}^{\infty} \text{Ad } e^{ith} \right) \stackrel{ec}{\sim} \alpha \otimes \gamma \otimes \left(\bigotimes_{n=1}^{\infty} \text{Ad } e^{ith} \right) \stackrel{ec}{\sim} \alpha \otimes \gamma \stackrel{ec}{\sim} \alpha.$$

Proposition 3.4 *If A is a UHF C^* -algebra such that $A \otimes A \cong A$, then the universal UHF flows on A are mutually almost conjugate. In particular a cocycle perturbation of a universal UHF flow γ is almost conjugate to γ .*

Proof. If α is a UHF flow and γ is a universal UHF flow, it follows that $\alpha^\infty \otimes \gamma^\infty \stackrel{ec}{\sim} \gamma$, where $\alpha^\infty = \bigotimes_{n=1}^{\infty} \alpha$ etc. Hence for any $\epsilon > 0$ there exists a flow ρ on A such that $\alpha^\infty \otimes \gamma^\infty \otimes \rho$ is conjugate to γ up to ϵ (i.e., $\|\alpha_t^\infty \otimes \gamma_t^\infty \otimes \rho_t - \varphi^{-1}\gamma_t\varphi\| < \epsilon$ for any $t \in [-1, 1]$ and for some isomorphism $\varphi : A^\infty \rightarrow A$). Hence $\alpha^\infty \otimes \gamma^\infty \otimes \rho = \alpha \otimes \alpha^\infty \otimes \gamma^\infty \otimes \rho$ is conjugate to $\alpha \otimes \gamma$ up to ϵ . Hence $\alpha \otimes \gamma \stackrel{ac}{\sim} \gamma$. If both α and γ are universal, it follows that $\alpha \stackrel{ac}{\sim} \alpha \otimes \gamma \stackrel{ac}{\sim} \gamma$.

Remark 3.5 Let A be a UHF C^* -algebra with $A \otimes A \cong A$. Then there is a *universal* automorphism γ of A in the sense that $\gamma \otimes \alpha$ is cocycle conjugate to γ for any automorphism α of A . Namely an automorphism with the Rohlin property is universal (see, e.g., [3, 11, 10, 7]). The universal automorphisms are mutually almost conjugate.

4 The flip automorphism

When A is a UHF C^* -algebra, the flip automorphism σ of $A \otimes A$ defined by $\sigma(x \otimes y) = y \otimes x$, $x, y \in A$ is approximately inner (since σ induces the identity map on $K_0(A \otimes A)$, which has rank one). Hence there is a sequence (u_n) of unitaries in $A \otimes A$ such that $\sigma = \lim_n \text{Ad } u_n$.

Let α be a flow on A . Since $\sigma(\alpha \otimes \alpha) = (\alpha \otimes \alpha)\sigma$, we can ask a question of whether one can put an extra condition, in connection with α , on the sequence (u_n) above.

Before turning to this special situation we present the following two propositions to clarify the conditions we are thinking of (cf. [8]).

Proposition 4.1 *Let A be a unital C^* -algebra and let α be a flow on A and $\sigma \in \text{Aut}(A \otimes A)$ such that $\sigma^{-1}\alpha\sigma = \alpha$. Then the following conditions are equivalent:*

1. *There is a sequence (u_n) of unitaries in $D(\delta_\alpha)$ such that $\sigma = \lim_n \text{Ad } u_n$ and $\sup_n \|\delta_\alpha(u_n)\| < \infty$.*

2. There is a sequence (u_n) of unitaries in A such that $\sigma = \lim_n \text{Ad } u_n$ and $(t \mapsto \alpha_t(u_n))$ are equi-continuous in n .

Proof. If (1) is satisfied, then so is (2) for the same (u_n) . To go from (2) to (1) we may need to modify (u_n) . Since a similar technique will apply in the proof of the following proposition, we do not give it here.

We will express the conditions in the above proposition by saying that σ is α -smoothly approximately inner, which is weaker than the condition that σ is α -invariantly approximately inner, which will appear in the following.

Proposition 4.2 *Let A be a unital C^* -algebra and let α be a flow on A and $\sigma \in \text{Aut}(A)$ such that $\sigma^{-1}\alpha\sigma = \alpha$. Then the following conditions are equivalent:*

1. There is a sequence (u_n) of unitaries in $D(\delta_\alpha)$ such that $\sigma = \lim_n \text{Ad } u_n$ and $\lim_n \|\delta_\alpha(u_n)\| = 0$.
2. There is a sequence (u_n) of unitaries in A such that $\sigma = \lim_n \text{Ad } u_n$ and $\|\alpha_t(u_n) - u_n\| \rightarrow 0$ uniformly in t on every compact subset of $\mathbf{R}$.
3. There is a sequence (u_n) of unitaries in A such that $\sigma = \lim_n \text{Ad } u_n$ and $\|\alpha_t(u_n) - u_n\| \rightarrow 0$ for every $t \in \mathbf{R}$.

Proof. It is obvious that (1) $\Rightarrow$ (2) $\Rightarrow$ (3).

(3) $\Rightarrow$ (2). Let $\epsilon > 0$ and define

$$I_n = \{t \in \mathbf{R} \mid \sup_{m \geq n} \|\alpha_t(u_m) - u_m\| \leq \epsilon\}.$$

Since $\cup_n I_n = \mathbf{R}$ and I_n 's are closed, $\cup_n I_n^\circ$ is dense in $\mathbf{R}$ by the Baire Category theorem, where I_n° is the interior of I_n . If $(a, a + \delta) \subset I_n^\circ$ for some $a \in \mathbf{R}$, $\delta > 0$, and n , then $\|\alpha_t(u_m) - u_m\| \leq 2\epsilon$ for $t \in (-\delta, \delta)$ and $m \geq n$. Then the rest is easy.

(2) $\Rightarrow$ (1). Let f be a C^∞ -function on $\mathbf{R}$ with compact support such that $f \geq 0$ and $\int f(t)dt = 1$ and let, for a small $\epsilon > 0$,

$$x_{n\epsilon} = \epsilon \int f(\epsilon t) \alpha_t(u_n) dt.$$

Since

$$\|x_{n\epsilon} - u_n\| \leq \epsilon \int f(\epsilon t) \|\alpha_t(u_n) - u_n\| dt,$$

one can choose a sequence (ϵ_n) such that $\epsilon_n \searrow 0$ and $\|y_n - u_n\| \rightarrow 0$ with $y_n = x_{n\epsilon_n}$. Note that $y_n \in D(\delta_\alpha)$ and that

$$\|\delta_\alpha(y_n)\| \leq \epsilon_n \int |f'(t)| dt.$$

If we denote by v_n the unitary obtained by the polar decomposition of y_n , the sequence (v_n) satisfies the desired properties.

If a flow α has more than one α -KMS states for some inverse temperature, the following shows that the flip is not α -smoothly approximately inner.

Proposition 4.3 *Let A be a UHF C^* -algebra and α a flow on A and suppose that the flip is α -smoothly approximately inner (or more precisely, $\alpha \otimes \alpha$ -smoothly approximately inner), i.e., the conditions in 4.1 are satisfied for $A \otimes A$, $\alpha \otimes \alpha$, and the flip automorphism σ in place of A , α , and σ respectively. Then the set of α -KMS states is a singleton (if not empty) for each inverse temperature.*

Proof. Let ω_i be an α -KMS state at inverse temperature $c \in \mathbf{R}$ for $i = 1, 2$. Then $\omega_1 \otimes \omega_2$ is an $\alpha \otimes \alpha$ -KMS state of $A \otimes A$ at c . Since the flip σ commutes with $\alpha \otimes \alpha$ and the flip is α -smoothly approximately inner, the flip leaves each $\alpha \otimes \alpha$ -KMS state invariant by the following lemma, i.e., $(\omega_1 \otimes \omega_2)\sigma = \omega_1 \otimes \omega_2$ or $\omega_2 \otimes \omega_1 = \omega_1 \otimes \omega_2$. Hence $\omega_1 = \omega_2$.

The following lemma is shown by M. Fannes et al [8] (or 5.3.33A of [4] vol. II), but we will present another proof based on the definition of the KMS condition in terms of holomorphic functions.

Lemma 4.4 *Let A be a unital C^* -algebra and let α be a flow and $\sigma \in \text{Aut}(A)$ such that $\sigma^{-1}\alpha\sigma = \alpha$. Suppose that σ is α -smoothly approximately inner. Then $\omega\sigma = \omega$ for any α -KMS states at inverse temperature $c \in \mathbf{R}$.*

Proof. Let ω be an α -KMS state at inverse temperature $c > 0$. (The case $c < 0$ entails just some notational changes and the case $c = 0$ is trivial.) By definition, for any $x, y \in A$ there is a bounded continuous function $F(z)$ on $S_c = \{z \in \mathbf{C} \mid 0 \leq \Im z \leq c\}$ such that F is analytic in the interior of S_c and

$$F(t) = \omega(x\alpha_t(y)), \quad F(t + ic) = \omega(\alpha_t(y)x)$$

for $t \in \mathbf{R}$. Let (u_n) be the sequence of unitaries as given in 4.1 and denote by $F_{n,x}$ the function F obtained by taking $u_n x$ for x and u_n^* for y . In particular $F_{n,x}$ satisfies

$$F_{n,x}(t) = \omega(u_n x \alpha_t(u_n^*)), \quad F_{n,x}(t + ic) = \omega(\alpha_t(u_n^*) u_n x),$$

for $t \in \mathbf{R}$. Since $t \mapsto \alpha_t(u_n^*) u_n$ is equi-continuous in n , we may assume by passing to a subsequence that $F_{n,y}$ converges, for $y = x$ and $y = 1$, uniformly on every compact subset of the boundary ∂S_c . Then $F_{n,y}$ converges, say to F_y , uniformly on every compact subset of S_c . It follows that F_y is a bounded continuous function on S_c which is analytic in the interior.

Suppose further that ω is factorial, which causes no loss of generality since the extreme KMS states are all factorial. Then since $(\alpha_t(u_n^*) u_n)$ is central, it follows that $\omega(\alpha_t(u_n^*) u_n x)$ converges to $F_1(t + ic)\omega(x)$. This implies that $F_x(z) = F_1(z)\omega(x)$ for $z \in \mathbf{R} + ic$ and hence for $z \in S_c$. Since $F_x(0) = \omega\sigma(x)$ and $F_1(0) = 1$, we obtain that $\omega\sigma(x) = \omega(x)$. Hence $\omega\sigma = \omega$.

We may apply the above proposition 4.3 to the one-dimensional quantum lattice systems (since the *bounded surface energy* condition obviously implies that the flip is α -smoothly approximately inner), where the uniqueness of KMS states is of course well-known (see [17]). This is yet another proof.

When L is a bounded linear map of a C^* -algebra B into a C^* -algebra A , we denote by $\|L\|_{cb}$ the completely bounded norm defined by $\sup_n \|L \otimes \text{id}_n\|$, where id_n is the identity map on the matrix algebra M_n so that $L \otimes \text{id}_n$ is a linear map of $B \otimes M_n$ into $A \otimes M_n$. In the following proposition we will use such a norm when B is finite-dimensional and L is a difference of homomorphisms. In this case it follows from Christensen's result [5] that for any $\epsilon > 0$ there is a $\delta > 0$ such that if $\|L\| < \delta$ then $\|L\|_{cb} < \epsilon$. (Because if $L = \phi_1 - \phi_2$ has sufficiently small norm with ϕ_i a unital homomorphism of B into A , then there is a unitary $u \in A$ such that $\|u - 1\|$ is small and $\phi_2 = \text{Ad } u \phi_1$, which entails that $\|L\|_{cb} = \|(\text{id} - \text{Ad } u)\phi_1\|_{cb} \leq 2\|u - 1\|$.)

Proposition 4.5 *Let A be a UHF C^* -algebra with $A \otimes A \cong A$ and α a flow on A . Suppose that there exists a sequence (φ_n) in $\text{Aut}(A)$ such that $\varphi_n^{-1} \alpha \varphi_n \rightarrow \alpha$ and $(\varphi_n(x))$ is central for any $x \in A$ and that the flip is α -invariantly approximately inner.*

Then for any unital finite type I subfactor A_1 of A and $\epsilon > 0$ there is a finite type I subfactor B of A with $B \supset A_1$ such that if $\varphi \in \text{Aut}(A)$ satisfies that for any $t \in I = [-1, 1]$,

$$\|(\alpha_t - \varphi^{-1} \alpha_t \varphi)|B\|_{cb} < \epsilon'$$

for some $\epsilon' > 0$, then there is a unitary $u \in A$ such that

$$\begin{aligned} \max_{t \in I} \|\alpha_t(u) - u\| &< \epsilon' + \epsilon, \\ \|(\varphi - \text{Ad } u)|A_1\| &< \epsilon. \end{aligned}$$

Proof. Let σ denote the flip of $A \otimes A$. By the assumption on α there exists a unitary $u \in A \otimes A$ such that

$$\begin{aligned} \text{Ad } u|A_1 \otimes A_1 &= \sigma|A_1 \otimes A_1, \\ \max_{t \in I} \|\alpha_t \otimes \alpha_t(u) - u\| &< \epsilon/2. \end{aligned}$$

We may suppose that there is a finite type I subfactor B of A such that $B \otimes B \ni u$ and $B \supset A_1$.

By using the (φ_n) and φ in the statement, we define linear maps $\iota \otimes \varphi_n$ and $\varphi \otimes \varphi_n$ of the algebraic tensor product $A \odot A$ into A by

$$\begin{aligned} \iota \otimes \varphi_n(x \otimes y) &= x\varphi_n(y), \\ \varphi \otimes \varphi_n(x \otimes y) &= \varphi(x)\varphi_n(y). \end{aligned}$$

Since $(\varphi_n(y))$ is a central sequence for any $y \in A$, both $\iota \otimes \varphi_n$ and $\varphi \otimes \varphi_n$ are *approximate* homomorphisms. Thus $\iota \otimes \varphi_n(u)$ and $\varphi \otimes \varphi_n(u)$ are close to unitaries for the $u \in B \otimes B$ above for all large n .

Since

$$\|\alpha_t(\iota \otimes \varphi_n)(u) - \iota \otimes \varphi_n(\alpha_t \otimes \alpha_t(u))\|$$

converges to zero uniformly in $t \in I$, we obtain that

$$\|\alpha_t(\iota \otimes \varphi_n)(u) - \iota \otimes \varphi_n(u)\| < \epsilon/2$$

for $t \in I$ and for all large n .

For a finite sum $\sum_i x_i \otimes y_i \in B \otimes B$ it follows that

$$\|\alpha_t(\varphi \otimes \varphi_n)(\sum_i x_i \otimes y_i) - (\varphi \otimes \varphi_n)(\sum_i \alpha_t(x_i) \otimes \alpha_t(y_i))\|$$

converges to

$$\|\sum_i \alpha_t \varphi(x_i) \otimes \alpha_t(y_i) - \sum_i \varphi \alpha_t(x_i) \otimes \alpha_t(y_i)\|$$

which is less than or equal to $\|(\alpha_t \varphi - \varphi \alpha_t)|B\|_{cb} \|\sum_i x_i \otimes y_i\|$. Hence it follows that

$$\|\alpha_t(\varphi \otimes \varphi_n)(u) - (\varphi \otimes \varphi_n)(\alpha_t \otimes \alpha_t(u))\| < \epsilon'$$

for any $t \in I$ and for all large n and that

$$\max_{t \in I} \|\alpha_t(\varphi \otimes \varphi_n)(u) - (\varphi \otimes \varphi_n)(u)\| < \epsilon' + \epsilon/2.$$

Thus we obtain that $\max_{t \in I} \|\alpha_t(y_n) - y_n\| < \epsilon' + \epsilon$ for $y_n = (\varphi \otimes \varphi_n)(u)(\iota \otimes \varphi_n)(u)$.

Since, for $x \in A_1$,

$$(\iota \otimes \varphi_n)(u)x \approx (\iota \otimes \varphi_n)(u \cdot x \otimes 1) = (\iota \otimes \varphi_n)(1 \otimes x \cdot u) \approx \varphi_n(x)(\iota \otimes \varphi_n)(u)$$

and

$$(\varphi \otimes \varphi_n)(u)\varphi_n(x) \approx (\varphi \otimes \varphi_n)(u \cdot 1 \otimes x) = (\varphi \otimes \varphi_n)(x \otimes 1 \cdot u) \approx \varphi(x)(\varphi \otimes \varphi_n)(u),$$

it follows that $\|y_n x - \varphi(x)y_n\| \rightarrow 0$ for any $x \in A_1$ as $n \rightarrow \infty$. Thus, by choosing a sufficiently large n and taking the unitary part of the polar decomposition of y_n , which is already close to a unitary, we obtain the conclusion. (As a matter of fact $\|(\text{Ad } u - \varphi)|A_1\|$ can be made small independently of ϵ .)

Theorem 4.6 *Let A be a UHF C^* -algebra with $A \otimes A \cong A$ and γ a universal UHF flow on A . If α is an approximately inner flow on A such that $\alpha \otimes \gamma \overset{c}{\approx} \alpha$, the following conditions are equivalent:*

1. $\alpha \overset{c}{\approx} \gamma$;
2. The flip is α -invariantly approximately inner.

Proof. (1) $\Rightarrow$ (2). We may assume that α is an inner perturbation of γ , i.e., $\delta_\alpha = \delta_\gamma + \text{ad } ih$ for some $h \in A_{sa}$. We know that there is a sequence (u_n) of unitaries in $D(\delta_\gamma) \odot D(\delta_\gamma)$ such that $u_n^* = u_n$, $\delta_{\gamma \otimes \gamma}(u_n) = 0$, and $\sigma = \lim_n \text{Ad } u_n$, where σ is the flip of $A \otimes A$. Hence it follows that $u_n \in D(\delta_{\alpha \otimes \alpha})$ and that

$$\delta_{\alpha \otimes \alpha}(u_n) = \text{ad}(ih \otimes 1 + 1 \otimes ih)(u_n) = (ih \otimes 1)u_n - u_n(1 \otimes ih) + (1 \otimes ih)u_n - u_n(ih \otimes 1)$$

converges to zero. This is what we wanted to show; see 4.2.

(2) $\Rightarrow$ (1). Since $\gamma^\infty \overset{c}{\sim} \gamma$, we see that there is a sequence (φ_n) in $\text{Aut}(A)$ such that $\varphi_n^{-1}\gamma\varphi_n \rightarrow \gamma$ and $(\varphi_n(x))$ is central for any $x \in A$. Since α is an approximately inner flow satisfying $\alpha \overset{c}{\sim} \alpha \otimes \gamma^\infty$ and any inner flow can be embedded into γ , we see that α also satisfies the above condition. Since the flip is both γ and α -invariantly approximately inner, we can apply 4.5 to both of them.

Let (A_n) be an increasing sequence of finite type I subfactors of A with $A = \overline{\bigcup_n A_n}$.

Let $\epsilon > 0$. For A_1 , α , and $2^{-1}\epsilon$ (in place of ϵ) we choose B_1 as B in 4.5. By 3.2 we choose $\varphi_1 \in \text{Aut}(A)$ such that for $t \in I = [-1, 1]$,

$$\|(\alpha_t - \varphi_1^{-1}\gamma_t\varphi_1)|B_1\|_{cb} < 2^{-2}\epsilon.$$

By slightly changing φ_1 if necessary we assume that $\varphi_1(B_1) \subset A_n$ for some $n > 1$; by passing to a subsequence of (A_n) we assume that $n = 2$; i.e., $\varphi_1(B_1) \subset A_2$. We then choose B_2 for A_2 , γ , and $2^{-2}\epsilon$ by 4.5 and choose $\varphi_2 \in \text{Aut}(A)$ by 3.3 such that for $t \in I$,

$$\|(\gamma_t - \varphi_2^{-1}\alpha_t\varphi_2)|B_2\|_{cb} < 2^{-3}\epsilon.$$

Here we again assume that $\varphi_2(B_2) \subset A_3$ as above. Since $B_2 \supset A_2 \supset \varphi_1(B_1)$, we have that

$$\|(\alpha_t - \varphi_1^{-1}\varphi_2^{-1}\alpha_t\varphi_2\varphi_1)|B_1\|_{cb} < (2^{-2} + 2^{-3})\epsilon.$$

By 4.5 we have a unitary $u_2 \in A_2$ such that

$$\begin{aligned} \|(\text{Ad } u_2\varphi_2\varphi_1 - \text{id})|A_1\| &< 2^{-1}\epsilon, \\ \max_{t \in I} \|\alpha_t(u_2) - u_2\| &< \epsilon. \end{aligned}$$

We choose B_3 for A_3 , α , and $2^{-3}\epsilon$ as in 4.5. We may further assume that $B_3 \ni u_2$. We then choose $\varphi_3 \in \text{Aut}(A)$ such that for $t \in I$,

$$\|(\alpha_t - \varphi_3^{-1}\gamma_t\varphi_3)|B_3\|_{cb} < 2^{-4}\epsilon.$$

Here again we assume that $\varphi_3(B_3) \subset A_4$. Since $\varphi_2(B_2) \subset A_3 \subset B_3$, we have that for $t \in I$,

$$\|(\gamma_t - \varphi_2^{-1}\varphi_3^{-1}\gamma_t\varphi_3\varphi_2)|B_2\|_{cb} < (2^{-3} + 2^{-4})\epsilon.$$

We then obtain a unitary $u_3 \in A$ such that

$$\begin{aligned} \|(\text{Ad } u_3\varphi_3\varphi_2 - \text{id})|A_2\| &< 2^{-2}\epsilon, \\ \max_{t \in I} \|\gamma_t(u_3) - u_3\| &< 2^{-1}\epsilon. \end{aligned}$$

We repeat this procedure.

With (φ_n) , (B_n) , and (u_n) obtained as above we proceed as follows. Let $u_1 = 1$, $v_1 = 1 = v_2$, and $v_3 = u_2$. We define a unitary v_n for $n \geq 4$ by

$$v_n = u_{n-1}\varphi_{n-1}(v_{n-1}^*).$$

Since $v_3 \in B_3$ and $\varphi_{n-1}(B_{n-1}) \subset B_n$, if $v_{n-1} \in B_n$, we have that $v_n \in B_n$ for all n . By letting $\phi_n = \text{Ad } u_n \varphi_n \text{Ad } v_n^*$, we obtain an almost commutative diagram (cf. [6]):

$$\begin{array}{ccccccc} A_1 & \xrightarrow{\text{id}} & A_3 & \xrightarrow{\text{id}} & A_5 & \longrightarrow & \dots \\ \phi_1 \downarrow & \phi_2 \nearrow & \phi_3 \downarrow & \phi_4 \nearrow & \downarrow & & \dots \\ A_2 & \xrightarrow{\text{id}} & A_4 & \xrightarrow{\text{id}} & A_6 & \longrightarrow & \dots \end{array}$$

This is almost commutative because

$$\begin{aligned} \phi_n \phi_{n-1} &= \text{Ad } u_n \varphi_n \text{Ad } (\varphi_{n-1}(v_{n-1})u_{n-1}^*) \text{Ad } u_{n-1} \varphi_{n-1} \text{Ad } v_{n-1}^* \\ &= \text{Ad } u_n \varphi_n \varphi_{n-1}, \end{aligned}$$

which is in the neighborhood of the inclusion $A_{n-1} \subset A_{n+1}$ of norm less than $2^{-n+1}\epsilon$. Since $(\phi_{2n+1}(x))$ is a Cauchy sequence for $x \in \cup_n A_n$, we obtain a homomorphism ϕ of A into A as the extension of the limit of (ϕ_{2n+1}) . Since the homomorphism defined as the limit of (ϕ_{2n}) is the inverse of ϕ , it follows that ϕ is an isomorphism of A onto A .

For $t \in I$ we compute:

$$\begin{aligned} &\|(\gamma_t \phi_{2n+1} - \phi_{2n+1} \alpha_t)|B_{2n+1}\| \\ &\leq 2\|\gamma_t(u_{2n+1}) - u_{2n+1}\| + \|(\gamma_t \varphi_{2n+1} \text{Ad } v_{2n+1}^* - \varphi_{2n+1} \text{Ad } v_{2n+1}^* \alpha_t)|B_{2n+1}\| \\ &\leq 2\|\gamma_t(u_{2n+1}) - u_{2n+1}\| + 2\|\alpha_t(v_{2n+1}^*) - v_{2n+1}^*\| + \|(\gamma_t \varphi_{2n+1} - \varphi_{2n+1} \alpha_t)|B_{2n+1}\| \\ &< 2^{-2n+2}\epsilon + 2^{-2n-2}\epsilon + 2\|\alpha_t(v_{2n+1}) - v_{2n+1}\|. \end{aligned}$$

Since, for $t \in I$,

$$\begin{aligned} \|\alpha_t(v_{2n+1}) - v_{2n+1}\| &\leq \|\alpha_t(u_{2n}) - u_{2n}\| + \|(\alpha_t \varphi_{2n} - \varphi_{2n} \gamma_t)|B_{2n}\| + \|\gamma_t(v_{2n}^*) - v_{2n}^*\| \\ &\leq 2^{-2n+2}\epsilon + 2^{-2n-1}\epsilon + \|\gamma_t(v_{2n}) - v_{2n}\|, \end{aligned}$$

and $\|\gamma_t(v_{2n}) - v_{2n}\| < 2^{-2n+3}\epsilon + 2^{-2n}\epsilon + \|\alpha_t(v_{2n-1}) - v_{2n-1}\|$ and since $v_2 = 1$, we have that for $t \in I$

$$\|\alpha_t(v_{2n+1}) - v_{2n+1}\| < 2\epsilon + 2^{-2}\epsilon < 3\epsilon.$$

From the above computations we have that for $t \in I$,

$$\|(\gamma_t \phi_{2n+1} - \phi_{2n+1} \alpha_t)|A_{2n+1}\| < 7\epsilon.$$

Thus it follows that for $t \in I$ and $x \in \cup_n A_n$,

$$\|\gamma_t \phi(x) - \phi \alpha_t(x)\| \leq 7\epsilon \|x\|.$$

This implies that α is almost conjugate to γ and hence $\alpha \overset{cc}{\approx} \gamma$.

When B_1 and B_2 are C^* -subalgebras of A and $\epsilon > 0$, we write $B_1 \overset{\epsilon}{\subset} B_2$ if for any $x_1 \in B_1$ there is an $x_2 \in B_2$ with $\|x_1 - x_2\| \leq \epsilon \|x_1\|$. We denote by $\text{dist}(B_1, B_2)$ the infimum of $\epsilon > 0$ such that $B_1 \overset{\epsilon}{\subset} B_2$ and $B_2 \overset{\epsilon}{\subset} B_1$.

Let A be a UHF C^* -algebra and α a flow on A . We say that α is a *quasi-UHF* flow if there exists a sequence (A_n) of finite type I subfactors of A such that $A = \overline{\bigcup_n A_n}$ and

$$\sup_{|t| \leq 1} \text{dist}(A_n, \alpha_t(A_n))$$

converges to zero as $n \rightarrow \infty$. Then from 6.5 of [5] it follows that

$$\sup_{|t| \leq 1} \text{dist}(A_n \otimes A_n, \alpha_t(A_n) \otimes \alpha_t(A_n))$$

converges to zero as $n \rightarrow \infty$.

The following is an attempt to prove that the quasi-UHF flows are UHF.

Corollary 4.7 *Let A be a UHF C^* -algebra such that $A \otimes A \cong A$ and γ a universal UHF flow on A . If α is an approximately inner, quasi-UHF flow on A , then $\alpha \otimes \gamma$ is a cocycle perturbation of a (universal) UHF flow.*

Proof. There exists a sequence (ϵ_n) with $\epsilon_n \searrow 0$ satisfying: For any $x \in A_n \otimes A_n$ with $\|x\| \leq 1$ and $t \in I = [-1, 1]$ there is an $x_t \in A_n \otimes A_n$ with $\|x_t\| \leq 1$ such that $\|\alpha_t \otimes \alpha_t(x) - x_t\| \leq \epsilon_n$.

Let u_n be a self-adjoint unitary in $A_n \otimes A_n$ which implements the flip $\sigma|_{A_n \otimes A_n}$.

Since, for $x \in A_n$ with $\|x\| \leq 1$,

$$(\alpha_t \otimes \alpha_t)(u_n)x \approx (\alpha_t \otimes \alpha_t)(u_n x_{-t}) = (\alpha_t \otimes \alpha_t)(\sigma(x_{-t})u_n) \approx \sigma(x)(\alpha_t \otimes \alpha_t)(u_n),$$

it follows that $\|[(\alpha_t \otimes \alpha_t)(u_n)u_n, x]\| \leq 2\epsilon_n\|x\|$ for $x \in A_n \otimes A_n$. Since there is a $v_{nt} \in A_n \otimes A_n$ such that $\|(\alpha_t \otimes \alpha_t)(u_n) - v_{nt}\| \leq \epsilon_n$, $\|v_{nt}\| \leq 1$, $v_{nt}^* = v_{nt}$, and $\sigma(v_{nt}) = v_{nt}$, we have that $v_{nt}u_n \in A_n \otimes A_n$ is self-adjoint and $\|[v_{nt}u_n, x]\| \leq 4\epsilon_n\|x\|$ for $x \in A_n \otimes A_n$. Since $\|(v_{nt}u_n)^2 - 1\| = \|v_{nt}^2 - 1\| \leq 2\epsilon_n$, $v_{nt}u_n$ must be close to 1 or -1 . As we may choose v_{nt} , $t \in I$ to be continuous in t with $v_{n0} = u_n$, we have that $v_{nt}u_n$ is close to 1, i.e., $\|v_{nt}u_n - 1\| \leq 6\epsilon_n$. Hence we get that $\|(\alpha_t \otimes \alpha_t)(u_n) - u_n\| \leq 7\epsilon_n$ for $t \in I$.

Since there is a sequence (w_n) of unitaries in $A \otimes A$ such that $\sigma = \lim \text{Ad } w_n$ and $\gamma_t \otimes \gamma_t(w_n) = w_n$, we have that the sequence $(u_n \otimes w_n)$ satisfies that $\sigma = \lim \text{Ad}(u_n \otimes w_n)$ and $\|(\alpha_t \otimes \alpha_t \otimes \gamma_t \otimes \gamma_t)(u_n \otimes w_n) - u_n \otimes w_n\| \rightarrow 0$ uniformly in $t \in I$, i.e., the flip is $\alpha \otimes \gamma$ -invariantly approximately inner. By the previous theorem we can conclude that $\alpha \otimes \gamma$ is a cocycle perturbation of γ .

5 The CAR algebra

When A is the CAR algebra (i.e., $A \cong M_{2^\infty}$), we can give a universal flow in a simple way.

Theorem 5.1 *Let (λ_n) be a sequence in $\mathbf{R}$ and define a UHF flow α on $A = M_{2^\infty}$ by*

$$\alpha_t = \bigotimes_{n=1}^{\infty} \text{Ad} \begin{pmatrix} e^{it\lambda_n} & 0 \\ 0 & 1 \end{pmatrix}.$$

Then α is a universal UHF flow on A if and only if there is a subsequence (λ_{n_k}) such that $\lim_k \lambda_{n_k} = 0$ and $\sum_k \lambda_{n_k}^2 = \infty$.

Proof. If α is a universal UHF flow, then the τ bottom marginal spectrum of α must be full, i.e., $\Gamma_{\tau,-}(\alpha) = \mathbf{R}_+$, where τ is the tracial state of A (see [12]). If there is an $\epsilon > 0$ such that $\sum_{n:|\lambda_n|<\epsilon} \lambda_n^2 < \infty$, then $\Gamma_{\tau,-}(\alpha) \subset [\epsilon, \infty)$ by 5.3 of [12]. Hence there must be a subsequence (λ_{n_k}) such that $\lim_k \lambda_{n_k} = 0$ and $\sum_k \lambda_{n_k}^2 = \infty$.

Conversely suppose that (λ_n) satisfies the required property. Let (μ_n) be another sequence in $\mathbf{R}$ and denote by β the UHF flow constructed as in the above statement. What we have shown in [12] is that $\alpha \otimes \beta \stackrel{\mathcal{C}}{\sim} \alpha$ for all such β . Now what we have to show is that $\alpha \otimes \beta \stackrel{\mathcal{C}}{\sim} \alpha$ for any UHF flow β . As in [12] this follows easily from the following lemma.

Lemma 5.2 *Let (λ_n) be a sequence in $\mathbf{R}$ as in the above theorem and let T be a finite subset of $\mathbf{R}$ of order 2^k . For any $\epsilon > 0$ there exist an $N \in \mathbf{N}$ with $N > k$, a partition P of the power set $\mathcal{P}_N$ of $\{1, 2, \dots, N\}$ into 2^{N-k} sets of order 2^k , and a map F of $\mathcal{P}_N$ onto T such that for any $S \in P$, $F|_S$ is bijective and*

$$|E(x) - E(y) - F(x) + F(y)| < \epsilon, \quad x, y \in S,$$

where $E(x) = \sum_{i \in x} \lambda_i$.

Proof. This lemma is shown in the case $k = 1$ in [12]. We will extend the proof there to cover the general case $k > 1$, at the same time using the result for the case $k = 1$.

We write the elements of T as $\mu_1, \mu_2, \dots, \mu_{2^k}$ in the increasing order. Since only the differences between μ_i 's matter, we assume that $\mu_{2^k} = -\mu_1$.

For a large N it is certainly not difficult to find a subset $\{x_1, x_2, \dots, x_{2^k}\}$ of $\mathcal{P}_N$ such that $(E(x_1), E(x_2), \dots, E(x_{2^k}))$ is almost equal to $(\mu_1, \mu_2, \dots, \mu_{2^k}) + C$ for some $C \in \mathbf{R}$ since $\{E(x) \mid x \in \mathcal{P}_N\}$ is densely distributed in a large interval. We have to cover $\mathcal{P}_N$ with the disjoint union of such sets $\{x_1, x_2, \dots, x_{2^k}\}$. Then the map $F : \mathcal{P}_N \rightarrow T$ is an obvious one.

Let $\mu_{0,i} = \mu_i$ and

$$\mu_{j,i} = \frac{1}{2}(\mu_{2^j(i-1)+1} + \mu_{2^j i}), \quad i = 1, 2, \dots, 2^{k-j},$$

for $j = 1, 2, \dots, k$. Since $\mu_{2^k} = -\mu_1$, we have that $\mu_{k,1} = 0$. For a fixed j , $(\mu_{j,i})$ is increasing in i and

$$\mu_{j,2i-1} < \mu_{j+1,i} < \mu_{j,2i}.$$

By slightly changing μ_i 's we may assume that all those $\mu_{j,i}$'s belong to $\mathbf{Z}\delta$ for some $\delta > 0$, which is smaller than ϵ . Furthermore, by using the result for $k = 1$, we may assume that all $\lambda_n = \delta$.

Fix a large N and define $E_n(x)$ for $x \in \mathcal{P}_N$ and $n \in \mathbf{N}$ with $n \leq N$ by

$$\begin{aligned} E_n(x) &= \frac{1}{2} \left(\sum_{i \leq n, i \in x} \lambda_i - \sum_{i \leq n, i \notin x} \lambda_i \right) \\ &= (\#\{i \in x \mid i \leq n\} - n/2)\delta. \end{aligned}$$

We say that x is *good* if there is a subsequence $n_1, n_2, \dots, n_k$ of even integers such that

$$E_{n_j}(x) = \mu_{k-j, i_j}$$

for some i_j ; in particular $E_{n_k}(x) = \mu_{i_k}$. We will make the smallest possible choice for each n_j inductively for $j = 1, 2, \dots, k$. Since $\mu_{k-j-1, 2i-1} < \mu_{k-j, i} < \mu_{k-j-1, 2i}$, this entails $i_{j+1} = 2i_j - 1$ or $2i_j$.

We claim that if x is good, then x can be matched with other $2^k - 1$ good elements of $\mathcal{P}_N$ in a unique way so that the resulting sequence $(x_1, x_2, \dots, x_{2^k})$ satisfies that $(E_{n_k}(x_1), E_{n_k}(x_2), \dots, E_{n_k}(x_{2^k}))$ exactly equals $(\mu_1, \mu_2, \dots, \mu_{2^k})$. Since

$$x_i \cap \{n_k + 1, n_k + 2, \dots, N\}$$

is independent of i by the construction, it follows that $(E(x_1), E(x_2), \dots, E(x_{2^k}))$ is equal to $(\mu_1, \mu_2, \dots, \mu_{2^k}) + C$ for some C , where $E(x_i) = E_N(x_i)$.

If x is good as above, we define $x' \in \mathcal{P}_N$ by

$$x' = x \triangle \{n_{k-1} + 1, n_{k-1} + 2, \dots, n_k\},$$

where $\triangle$ denotes difference of sets. Then x' is again good; $E_{n_j}(x') = \mu_{k-j, i_j}$ for $j = 1, 2, \dots, k-1$ and $E_{n_k}(x') = \mu_\ell$, where $\ell = i_k + 1$ if i_k is odd and otherwise $\ell = i_k - 1$. In general if (μ_{k-j, i_j}) is monotone for $j = m, m+1, \dots, k$, we define $x' \in \mathcal{P}_N$ by

$$x' = x \triangle \{n_m + 1, n_m + 2, \dots, n_k\}.$$

Then we have that x' is good and $E_{n_k}(x') = \mu_\ell$, where $\ell = 2^{k-m}i_m$ if i_k is odd and otherwise $\ell = 2^{k-m}(i_m - 1) + 1$ (since $\mu_{k-m, i_m} = (\mu_{2^{k-m}(i_m-1)+1} + \mu_{2^{k-m}i_m})/2$ and i_k is either $2^{k-m}(i_m - 1) + 1$ or $2^{k-m}i_m$ by the monotonicity). If we make the smallest possible choice of n'_j such that $E_{n'_j}(x') = \mu_{k-j, i'_j}$, we have that $n'_k = n_k$, where the latter is the choice for x . We may call this process the *reflection* at (m, i_m) or loosely at μ_{k-m, i_m} . By applying the above process inductively we can obtain the other $2^k - 1$ good elements from one good x , which form the desired set of order 2^k . (To understand why we get a set

of 2^k elements in this way we should visualize a binary tree of depth k with each node labeled by μ_{ji} such that the root is labeled by $\mu_{k1} = 0$ and if a node is labeled by $\mu_{k-j,i}$ with $j < k$ or more precisely is addressed by $(k-j, i)$, meaning that it is at a distance $k-j$ from the root and at the i 'th position from the left among the nodes of distance $k-j$ from the root, then it has two children labeled by $\mu_{k-j-1,2i-1}$ and $\mu_{k-j-1,2i}$ from left to right; so the leaves are labeled by $\mu_1 = \mu_{0,1}, \mu_{0,2}, \dots, \mu_{0,2^k}$ from left to right. Any good element $x \in \mathcal{P}_N$ corresponds to a path from the root to a leaf of this binary tree, which is the path determined by $(k-j, i_j)$, $j = 0, 1, \dots, k$ with $i_0 = 1$ and (i_j) as given in the definition of *good*. By the procedure indicated above we get $2^k - 1$ elements corresponding to other $2^k - 1$ paths. Note that to each interior node there is a reflection to be applied; and there are $2^k - 1$ of them.)

In this way we obtain the family F_N of such sets of order 2^k . Since the procedure is canonical, F_N is a disjoint family. Let G_N be the union of elements of F_N . If $x \in \mathcal{P}_N \setminus G_N$, then x is not good and in particular $|E_n(x)| < \mu_{2^k}$ for all $n = 1, 2, \dots, N$.

For each $k \in \mathbf{Z}$ with $|k| < \mu_{2^k}/\delta \equiv K$, we have that

$$\frac{\#\{x \in \mathcal{P}_N \setminus G_N \mid E(x) = k\delta\}}{\#\{x \in G_N \mid E(x) = k\delta\}}$$

converges to zero as even N goes to infinity. Furthermore we have that

$$\frac{\#(\mathcal{P}_N \setminus G_N)}{\min_{|k| \leq K} \#\{x \in G_N \mid E(x) = k\delta\}}$$

converges to zero as even N goes to infinity (since the ratios among $\#\{x \in G_N \mid E(x) = k\delta\}$ for various k converge to 1).

For each $x \in \mathcal{P}_N \setminus G_N$ we specify a subset S_x of G_N of order $2K+1$ such that

$$\{E(y) \mid y \in S_x\} = \{k\delta \mid |k| \leq K\}.$$

If N is a sufficiently large even integer, we can specify S_x , $x \in \mathcal{P}_N \setminus G_N$ such that S_x 's are mutually disjoint.

Let $\{x_1, x_2, \dots, x_{2^k}\}$ be a subset of $\mathcal{P}_N \setminus G_N$. If $E(x_1) = k\delta$, we make the following substitutions (simultaneously):

$$y_{k-1} \leftarrow x_1, y_{k-2} \leftarrow y_{k-1}, \dots, y_{-K} \leftarrow y_{-K+1}, x_1 \leftarrow y_{-K},$$

where $y_k \in S_{x_1}$ satisfies $E(y_k) = k\delta$. After these substitutions we have that $E(x_1) = -K\delta = \mu_1$. By making suitable substitutions among $\{x_i\} \cup S_{x_i}$ for $i = 2, \dots, 2^k$ we have that $E(x_i) = \mu_i$, i.e., $\{x_1, x_2, \dots, x_{2^k}\}$ satisfies the desired property. By these substitutions we have introduced only an error of δ to some sets belonging to F_N , i.e., for $S \in F_N$ modified by these processes, the elements of S will be ordered as $x_1, x_2, \dots, x_{2^k}$ so that $|E(x_i) - \mu_i - C| = 0, \pm\delta$ for all i and for some C . This completes the proof.

References

- [1] O. Bratteli, *Derivations, dissipations and group actions on C^* -algebras*, Lecture Notes in Math. 1229, Springer, 1986.
- [2] O. Bratteli and A. Kishimoto, AF flows and continuous symmetries, preprint.
- [3] O. Bratteli, A. Kishimoto, M. Rørdam, and E. Størmer, The crossed product of a UHF algebra by a shift, *Ergod. Th. & Dynam. Sys.* 13 (1993), 615–626.
- [4] O. Bratteli and D.W. Robinson, *Operator algebras and quantum statistical mechanics, I, II*, Springer, 1979, 1996.
- [5] E. Christensen, Near inclusions of C^* -algebras, *Acta Math.* 144 (1980), 249–265.
- [6] G.A. Elliott, On the classification of C^* -algebras of real rank zero, *J. reine angew. Math.* 443 (1993), 179–219.
- [7] D.E. Evans and A. Kishimoto, Trace-scaling automorphisms of certain stable AF algebras, *Hokkaido Math. J.* 26 (1997), 211–224.
- [8] M. Fannes, P. Vanheuverzwijn, and A. Verbeure, Quantum energy-entropy inequalities: A new method for proving the absence of symmetry breaking, *J. Math. Phys.* 25 (1984), 76–78.
- [9] A. Kishimoto, Outer automorphism subgroups of a compact abelian ergodic action, *J. Operator Theory* 20 (1988), 59–67.
- [10] A. Kishimoto, The Rohlin property for automorphisms of UHF algebras, *J. reine angew. Math.* 465 (1995), 183–196.
- [11] A. Kishimoto, The Rohlin property for shifts on UHF algebras and automorphisms of Cuntz algebras, *J. Funct. Anal.* 140 (1996), 100–123.
- [12] A. Kishimoto, Locally representable one-parameter automorphism groups of AF algebras and KMS states, *Rep. Math. Phys.* 45 (2000), 333–356.
- [13] A. Kishimoto, Examples of one-parameter automorphism groups of UHF algebras, to appear in *Commun. Math. Phys.*
- [14] A. Kishimoto, Non-commutative shifts and crossed products, preprint.
- [15] H. Lin, Classification of simple C^* -algebra of tracial topological rank zero, preprint.
- [16] D. Olsen and G.K. Pedersen, Applications of the Connes spectrum to C^* -dynamical systems, III, *J. Funct. Anal.* 45, 357–390 (1982).
- [17] S. Sakai, *Operator Algebras in Dynamical Systems*, Cambridge Univ. Press, 1991.
- [18] S. Sakai, *C^* -algebras and W^* -algebras*, Classics in Math. Springer, 1998.